JOURNAL OF CHROMATOGRAPHY A

VOL. 771, NOS. 1 + 2 30 MAY 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Application of sequential paired covariance to liquid chromatography–mass spectrometry data. Enhancements in both the signal-to-noise ratio and the resolution of analyte peaks in the chromatogram
by D.C. Muddiman, B.M. Huang, G.A. Anderson, A. Rockwood, S.A. Hofstadler, M.S. Weir-Lipton, A. Proctor, Q. Wu and R.D. Smith (Richland, WA, USA) (Received 7 November 1996) 1

Productivity and operating regimes in protein chromatography using low-molecular-mass displacers
by S.R. Gallant and S.M. Cramer (Troy, NY, USA) (Received 18 November 1996) 9

Alkanesulfonate homologous series as eluent components in anion chromatography. Novel eluent selection system for monovalent analytes
by P. Hajós and G. Révész (Veszprém, Hungary) (Received 31 December 1996) 23

Economic approach to robotic sample pretreatment in high-performance liquid chromatography
by A. Tolokán, L. Gödörházy, V. Horváth, A. Hrabéczy-Páll, Z. Niegreisz, E. Kocsi, G. Horvai, I. Klebovich and K. Balogh-Nemes (Budapest, Hungary) (Received 18 February 1997) 35

Silicalite as a sorbent for solid-phase extraction
by D.L. Mayer and J.S. Fritz (Ames, IA, USA) (Received 3 December 1996) 45

Chemically modified polymeric resin used as sorbent in a solid-phase extraction process to determine phenolic compounds in water
by N. Masqué, M. Galià, R.M. Marcé and F. Borrull (Tarragona, Spain) (Received 23 January 1997) 55

Purification of *p*-nitrobenzyl C-functionalized diethylenetriamine pentaacetic acids for clinical applications using anion-exchange chromatography
by M.W. Brechbiel, P.M. Beitzel and O.A. Gansow (Bethesda, MD, USA) (Received 27 January 1997) 63

Preparative separation of bitter acids from hop extracts by centrifugal partition chromatography
by A.C.J. Hermans-Lokkerbol, A.C. Hoek and R. Verpoorte (Leiden, The Netherlands) (Received 23 December 1996) 71

Peptide separation by pH-zone-refining countercurrent chromatography
by Y. Ma and Y. Ito (Bethesda, MD, USA) (Received 24 December 1996) 81

Separation of cyclic nitroxide free radicals and their redox forms with dual microelectrochemical detection
by B.L. Whited, P.N. Motsegood and J.E. Baur (Normal, IL, USA) (Received 3 January 1997) 89

High-performance liquid chromatographic analysis of *Pseudomonas aeruginosa* phenazines
by R.O. Fernández and R.A. Pizarro (Buenos Aires, Argentina) (Received 21 January 1997) 99

Relationship between the hydrophobicity and specific hydrophobic surface area of pesticides determined by high-performance liquid chromatography compared with reversed-phase thin-layer chromatography
by T. Cserháti and E. Forgács (Budapest, Hungary) (Received 7 January 1997) 105

(Continued overleaf)

Contents (continued)

Qualitative and quantitative analysis of styrene and its various oligomers by liquid chromatography
by H. El Mansouri, N. Yagoubi and D. Ferrier (Chatenay-Malabry, France) (Received 23 January 1997) 111

Drug quantitation on a benchtop liquid chromatography–tandem mass spectrometry system
by P.R. Tiller, J. Cunniff, A.P. Land, J. Schwartz, I. Jardine, M. Wakefield and L. Lopez (San Jose, CA, USA) and J.F. Newton, R.D. Burton, B.M. Folk, D.L. Buhrman, P. Price and D. Wu (Malvern, PA, USA) (Received 8 February 1997) 119

Determination of cyanocobalamin in foods by high-performance liquid chromatography with visible detection after solid-phase extraction and membrane filtration for the precolumn separation of lipophilic species
by H. Iwase and I. Ono (Kawasaki, Japan) (Received 27 January 1997) 127

Separation and determination of four artificial sweeteners and citric acid by high-performance anion-exchange chromatography
by Q.-c. Chen, S.-f. Mou and K.-n. Liu, Z.-y. Yang and Z.-m. Ni (Beijing, China) (Received 23 December 1996) 135

Separation and determination of oligomers and homologues of aliphatic alcohol ethoxylates in textile lubricants and lubricant emulsion by high-performance liquid chromatography
by C. Sun and M. Baird (Galashiels, UK), H.A. Anderson (Aberdeen, UK) and D.L. Brydon (Galashiels, UK) (Received 24 January 1997) 145

Analysis of amine-containing phosphonates in detergent powders by anion-exchange chromatography with pulsed amperometric detection
by K. Mahabir-Jagessar Tewari and S.A.M. van Stroe-Bieze (Vlaardingen, Netherlands) (Received 7 February 1997) 155

Optimising the determination of nitrate and phosphate in sea water with ion chromatography using experimental design
by I. Dahllöf, O. Svensson and C. Torstensson (Göteborg, Sweden) (Received 10 February 1997) 163

Gas Chromatography

Separation of extracts from biological tissues into polycyclic aromatic hydrocarbon, polychlorinated biphenyl and polychlorinated dibenzo-*p*-dioxin/polychlorinated dibenzofuran fractions prior to analysis
by P.W. O'Keefe, J. Miller, R. Smith, S. Connor, W. Clayton and R. Storm (Albany, NY, USA) (Received 3 February 1997) . . 169

Analysis of chlorobenzenes in soils by headspace solid-phase microextraction and gas chromatography–ion trap mass spectrometry
by F.J. Santos, M.N. Sarrión and M.T. Galceran (Barcelona, Spain) (Received 30 January 1997) 181

Gas–liquid chromatography–mass spectrometry of primary and secondary fatty alcohols and diols as their *tert.*-butyldimethylsilyl derivatives
by D.L. Chance, K.O. Gerhardt and T.P. Mawhinney (Columbia, MO, USA) (Received 31 December 1996) 191

Analysis of phenolic compounds in effluent by solid-phase extraction and gas chromatography–mass spectrometry with direct on-column benzylation: sensitive negative ion chemical ionisation gas chromatography–mass spectrometry detection of phenyl benzyl ethers
by J. Cheung and R.J. Wells (Pymble, Australia) (Received 4 February 1997) 203

Prediction of the gas chromatographic relative retention times of flavonoids from molecular structure
by P. Payares, D. Díaz, J. Olivero and R. Vivas (Cartagena, Colombia) and I. Gómez (Granada, Spain) (Received 23 January 1997) 213

Determination of pesticides in environmental waters by solid-phase extraction and gas chromatography with electron-capture and mass spectrometry detection
by C. Aguilar, F. Borrull and R.M. Marcé (Tarragona, Spain) (Received 22 January 1997) 221

Determination of toluene-2,4-diisocyanate in environmental and workplace air by sampling onto Tenax-TA followed by thermal desorption and capillary gas chromatography using flame ionisation and ion-trap detection
by A.P. Bianchi and T.K.B. Joyner (Southampton, UK) (Received 21 January 1997) 233

Supercritical Fluid Chromatography

Retention of some components in supercritical fluid chromatography and application to bergamot peel oil fractionation
by P. Subra and A. Vega (Villetaneuse, France) (Received 29 January 1997) 241

Planar Chromatography

Application of a colour analyzer in quantitative thin-layer chromatography
by M. Petrović, M. Kaštelan-Macan, S. Andrašić and L. Bokić (Zagreb, Croatia) (Received 11 November 1996) 251

Contents (continued)

Electrophoresis

Enantioseparation in capillary electrophoresis using 2-hydroxypropyltrimethylammonium salt of β-cyclodextrin as a chiral selector
by G. Schulte, B. Chankvetadze and G. Blaschke (Münster, Germany) (Received 20 January 1997) 259

Determination of cinnamaldehyde, cinnamic acid, paeoniflorin, glycyrrhizin and [6]-gingerol in the traditional Chinese medicinal preparation Kuei-chih-tang by cyclodextrin-modified micellar electrokinetic chromatography
by H.-Ya Huang, K.-L. Kuo and Y.-Z. Hsieh (Hsinchu, Taiwan, R.O.C) (Received 5 February 1997) 267

Non-aqueous capillary electrophoretic separation of Brønsted acids as heteroconjugated anions
by T. Okada (Tokyo, Japan) (Received 21 January 1997) 275

Determination of oxalate in parenteral nutrition solutions by capillary electrophoresis
by B.C. Nelson, P.C. Uden, G.F. Rockwell, K.M. Gorski and Z.B. Aguilera (Amherst, MA, USA) (Received 28 January 1997) 285

Determination of long-chained fatty acids using non-aqueous capillary electrophoresis and indirect UV detection
by E. Drange and E. Lundanes (Oslo, Norway) (Received 23 January 1997) 301

Optical resolution of phenylthiohydantoin-amino acids by capillary electrophoresis for protein sequencing
by Y. Kurosu, K. Murayama, N. Shindo, Y. Shisa, Y. Satou and N. Ishioka (Tokyo, Japan) (Received 3 February 1997) 311

Analysis of DNA fragmentation using a dynamic size-sieving polymer solution in capillary electrophoresis
by B.A. Siles, Z.E. Nackerdien and G. Bruce Collier (Gaithersburg, MD, USA) (Received 14 January 1997) 319

Wavelength-resolved laser-induced fluorescence detection in capillary electrophoresis: naphthalenesulphonates in river water
by S.J. Kok, E.M. Kristenson, C. Gooijer, N.H. Velthorst and U.A.Th. Brinkman (Amsterdam, Netherlands) (Received 29 January 1997) 331

SHORT COMMUNICATIONS

Column Liquid Chromatography

Chiral resolution of 4-substituted pyrrolidin-2-ones using high-performance liquid chromatography on cellulose and amylose chiral stationary phases
by C. Vaccher, M.-P. Vaccher and J.-P. Bonte (Lille, France) (Received 6 November 1996) 343

Determination of furazolidone in animal feeds using liquid chromatography with UV and thermospray mass spectrometric detection
by R.J. McCracken and D.G. Kennedy (Belfast, UK) (Received 6 February 1997) 349

Ion chromatographic procedures for analysis of total fluoride content in dentifrices
by T.A. Biemer, N. Asral and A. Sippy (Morris Plains, NJ, USA) (Received 10 January 1997) 355

Column Liquid and Gas Chromatography

Determination of morphine and its analogues in urine by on-line coupled reversed-phase liquid chromatography–gas chromatography with on-line derivatization
by T. Hyötyläinen, H. Keski-Hynnilä and M.-L. Riekkola (Helsinki, Finland) (Received 30 January 1997) 360

Identification of potential migrants from a vinylic organosol varnish by gas chromatography–mass spectrometry and liquid chromatography–mass spectrometry
by S. Cottier and A.M. Riquet (Jouy en Josas, France), A. Feigenbaum (Reims, France), B. Pollet and C. Lapierre (Thiverval-Grignon, France) and P. Mortreuil (Voreppe, France) (Received 30 January 1997) 366

Electrophoresis

Comparisons of the mobilities of salt ions obtained by the moving boundary method and two empirical equations in capillary electrophoresis
by C.-X. Cao (Wuhu Anhui, China) (Received 6 January 1997) 374

Off-column chemiluminescence detection in capillary electrophoresis
by Y.-T. Lee and C.-W. Whang (Taichung, Taiwan) (Received 7 January 1997) 379

Contents (continued)

LETTER TO THE EDITOR 385

ERRATA

Erratum to "Reversed-phase high-performance liquid chromatographic determination of ginsenosides of *Panax quinquefolium*". [J. Chromatogr. A, 755 (1996) 11] by W.A. Court, J.G. Hendel and J. Elmi (Delhi, Canada) (Received 10 March 1997) 387

Erratum to 'Polymerase chain reaction heteroduplex polymorphism analysis by entangled solution capillary electrophoresis'. [J. Chromatogr. A, 677 (1994) 169–177] by J. Cheng, T. Kasuga, K. R. Mitchelson and E.R.T. Lightly (Aberdeen, UK), N.D. Watson (Glasgow, UK) and W.J. Martin and D. Atkinson (Aberdeen, UK) (Received 21 February 1997) 389

AUTHOR INDEX VOL. 771 391

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc